

Vascular Function

Hiroaki Shimokawa, Kimio Satoh

Vascular function is regulated by many cell components, including endothelial cells (ECs), vascular smooth muscle cells (VSMCs), and adventitial tissues with inflammatory cells, autonomic nervous system, and vasa vasorum. The interactions among these cells/tissues are substantially involved in the vascular health and disease. All these cell components secrete several growth factors, cytokines/chemokines, and extracellular matrix, all of which contribute to vascular homeostasis. Among them, the growth factors secreted from VSMCs play important roles in the vascular remodeling process because they mediate various cellular responses. In addition, oxidative stress is one of the important stimuli that modulate VSMC function and proliferation. Many recent publications in *ATVB* have added further information on the roles of each cell component in the regulation of vascular function and diseases, including cytokines/chemokines, matrix metalloproteinases (MMPs), VSMC proliferation, and inflammation. Some publications have also provided a novel role of perivascular adipose tissues and perivascular stem cells that contribute to the development of vascular diseases. This article will focus on these recent publications in *ATVB*.

ECs for Vascular Function

ECs are exposed to blood flow, contributing to the maintenance of vessel structure and preservation of vascular functions. However, the effects of hemodynamic forces on cell signaling in the endothelium are still obscure. Several recent articles in *ATVB* addressed the role of ECs in the vascular function and mechanistic links to vascular diseases. Walshe et al¹ demonstrated that transforming growth factor- β (TGF- β) signaling mediates the protective effects of shear stress on ECs. They examined the potential role of TGF- β signaling in mediating the protective effects of shear stress on ECs. Finally, they concluded that shear stress induces TGF- β 3 signaling and subsequent activation of Kruppel-like factor 2 and nitric oxide (NO), demonstrating a novel role for TGF- β 3 in the maintenance of endothelial homeostasis in a hemodynamic environment.¹ This article provides a novel effect of shear stress on ECs in which induction of TGF- β 3 expression leads to the activation of Kruppel-like factor 2 and NO signaling. Approaches designed to assess endothelial functions, as well as gene expressions in vivo, provide a novel finding in

the analyses of cardiovascular diseases. As Enkhjargal et al² have recently demonstrated, endothelial AMP-activated protein kinase mediates endothelium-dependent hyperpolarization responses and blood pressure regulation in mice. ECs are central to the initiation of vascular diseases, yet there has been limited success in studying their gene expressions in the arteries in vivo. To address this, Erbilgin et al³ developed a novel method for determining the global transcriptional changes that occur in the mouse endothelium in response to atherogenic conditions and applied it to investigate inflammatory stimuli. They demonstrated that RNA can be isolated from mouse aortic ECs after in vivo and ex vivo treatments. In addition, they applied this method to identify a group of novel causative genes in vascular diseases. Choi et al⁴ demonstrated that globotriaosylceramide accelerates endocytosis and lysosomal degradation of endothelial K_{Ca}3.1 via a clathrin-dependent process, leading to endothelial dysfunction in Fabry disease. In this study, the authors demonstrated that glycosphingolipid globotriaosylceramide modulates K_{Ca}3.1 expression in ECs via clathrin-dependent and early endosome antigen 1-enriched endosome-mediated lysosomal degradation. Specifically, inhibition of globotriaosylceramide -induced K_{Ca}3.1 degradation through clathrin knockdown suggested that the K_{Ca}3.1 protein is internalized by globotriaosylceramide via a clathrin-dependent process. On the basis of the study, the authors proposed a novel mechanism underlying globotriaosylceramide-mediated EC dysfunction, which is associated with suppression of endothelium-derived hyperpolarizing factors.^{4,5}

This finding is particularly important in the clinical setting. H₂O₂ plays a crucial role as a signaling molecule at physiological low concentrations, which is one of the endothelium-dependent hyperpolarization factors that modulate vascular tone, especially in microvessels and in human coronary arteries. Thus, physiological levels of reactive oxygen species (ROS) levels are important for the regulation of EC functions. Although high levels of ROS induce vascular dysfunction, strictly controlled ROS formation mediates several important physiological vascular functions. However, it has long been awaited a plausible explanation and mechanisms for the biphasic effect of ROS on vascular function. A recent article in *ATVB* by Wood et al⁶ demonstrated that eNOS in circulating erythrocytes also contributes to the regulation of systemic blood pressure and vascular homeostasis. Moreover, Ciccarelli et al⁷ demonstrated that endothelial G-protein-coupled receptor kinase 2 removal induces endothelial dysfunction by increasing endothelial ROS production. Furthermore, Zippel et al⁸ demonstrated that transforming growth factor- β -activated kinase 1 regulates redox balance in ECs by AMP-activated protein kinase α 1, which acts as a metabolic master switch regulating several intracellular systems. In addition, Iso et al⁹ clearly demonstrated that capillary endothelial fatty acid-binding protein 4/5 is required for fatty

From the Department of Cardiovascular Medicine, Tohoku University Graduate School of Medicine, Sendai, Japan.

Correspondence to Hiroaki Shimokawa, MD, PhD, Department of Cardiovascular Medicine, Tohoku University Graduate School of Medicine, Sendai 980-8574, Japan. E-mail shimo@cardio.med.tohoku.ac.jp (*Arterioscler Thromb Vasc Biol.* 2014;34:2359-2362.)

© 2014 American Heart Association, Inc.

Arterioscler Thromb Vasc Biol is available at <http://atvb.ahajournals.org>
DOI: 10.1161/ATVBAHA.114.304119

acid transport into fatty acid-consuming tissues, including the heart, which controls the metabolism of energy substrates in fatty acid-consuming organs. All these recent reports focus on the crucial role of redox balance in the regulation of vascular function.

Oxidative stress causes vascular dysfunction, which is closely associated with vascular inflammation. Inflammation plays a crucial role in the regulation of endothelial functions and angiogenesis. A recent article in *ATVB* by Wagner et al¹⁰ examined the capacity of toll-like receptor 2–blocking Abs to modulate angiogenic responses of ECs in vitro and in vivo. They identified a novel molecular mechanism linking toll-like receptor 2 to angiogenic processes that is independent from the activation of inflammatory cascades, further supporting the concept of a beneficial effect of toll-like receptor 2 inhibition for EC function in vascular disease.¹⁰ Plasma levels of high-density lipoprotein (HDL) cholesterol are inversely associated with the risk of cardiovascular diseases. Multiple lines of evidence indicate that salutary effects of HDL on ECs include antioxidative, anti-inflammatory, antithrombotic, and proangiogenic effects. A recent article in *ATVB* by Tatematsu et al¹¹ reported that endothelial lipase acts on HDL to promote activation of S1P₁ (sphingosine-1-phosphate receptor 1). They also showed that endothelial lipase is a key player in facilitating the activation of HDL-dependent signals that lead to S1P₁ receptor activation, as well as Akt and eNOS phosphorylation.¹¹ These results provide new insights into the molecular mechanisms through which HDL, endothelial lipase, and their interaction could modulate angiogenesis and vascular function.¹¹

VSMCs for Vascular Function

Oxidative stress plays a major role in the pathogenesis of vascular diseases. However, it remains to be elucidated which ROS promotes the development of vascular disease. A recent article in *ATVB* by Parastatidis et al¹² examined the effect of hydrogen peroxide (H₂O₂)–degrading enzyme catalase on the formation of aortic aneurysms. They showed that restoration of catalase activity in the vascular wall enhances aortic VSMC survival and prevents aneurysm formation primarily through modulation of MMP activity.¹² They concluded that altered H₂O₂ signaling as a consequence of decreased catalase activity in the vasculature could be an early insult that leads to aortic dilatation.¹²

About the role of ROS for the development of vascular diseases, Soe et al¹³ also demonstrated that cyclophilin A plays a crucial role in the translocation of Nox enzymes, such as p47phox, which are known to contribute to VSMC proliferation and development of vascular diseases. Because ROS production by Nox enzymes activates other oxidase systems, cyclophilin A and Nox enzymes amplify ROS formation in a synergistic manner, leading to increased oxidative stress. Further knowledge of the role of cyclophilin A signaling on vascular cell components will contribute to the development of novel therapies for cardiovascular diseases. Rho-kinase is an important downstream effector of the small GTP-binding protein Rho and plays a crucial role in migration and proliferation of VSMCs. The RhoA/Rho-kinase pathway plays an important role in contraction, motility, and proliferation of

VSMCs, leading to the development of vascular diseases. The expression of Rho-kinase is accelerated by inflammatory stimuli, which upregulates nicotinamide adenine dinucleotide phosphate oxidase and further augments ROS formation. There are 2 isoforms of Rho-kinase, ROCK1 and ROCK2, and they have different functions. However, specific roles and functional differences between ROCK1 and ROCK2 remain to be elucidated in VSMCs. In a recent issue of *ATVB*, Shimizu et al¹⁴ demonstrated that ROCK2 in VSMCs contributes to VSMC proliferation and pulmonary hypertension in mice and in humans. Furthermore, Ikeda et al¹⁵ demonstrated that the Rho-kinase pathway (especially ROCK2) plays a crucial role in pressure-overload–induced right ventricular hypertrophy and dysfunction. Because secretion is regulated by Rho-kinase activity, the cyclophilin A/Rho-kinase pathway contributes to augmentation of oxidative stress and the development of cardiovascular diseases.

MMPs modulate extracellular matrix and regulates migration and proliferation of VSMCs. In a recent issue of *ATVB*, Xiao et al¹⁶ demonstrated that MMP-8 promotes VSMC proliferation and neointima formation. In this study, the authors clearly demonstrated that MMP-8 enhances VSMC proliferation via an ADAM10, N-cadherin, and β -catenin–mediated pathway and plays an important role in neointima formation. Integrins also contribute to vascular morphogenesis through regulation of adhesion and assembly of the extracellular matrix. Turlo et al¹⁷ showed an essential function of β 1-integrin in the maintenance of vasomotor control and highlighted a critical role for β 1-integrin in VSMC survival. Furthermore, Sutliff et al¹⁸ demonstrated that Poldip2 knock-down reduces H₂O₂ production in vivo, leading to increases in extracellular matrix, greater vascular stiffness, and impaired agonist-mediated contraction. Abnormal proliferation and migration of VSMCs are critical events in the progression of several vascular diseases, where AMP-activated protein kinase plays a crucial role in cellular proliferation and migration. In a recent issue of *ATVB*, Song et al¹⁹ demonstrated that deletion of AMP-activated protein kinase α 2 causes aberrant VSMC migration with accelerated neointima formation in vivo. In the pathogenesis of pulmonary hypertension, MMP activation and VSMC proliferation/migration are substantially involved. Revuelta-López et al²⁰ showed a crucial role of low-density lipoprotein receptor–related protein 1–mediated Pyk2 phosphorylation on hypoxia-induced MMP-9 activation, VSMC migration, and hypoxia-induced pulmonary hypertension. Low-density lipoprotein receptor–related protein 1 is a large endocytic and signaling receptor that is abundant in VSMCs. Muratoglu et al²¹ also suggested a critical role for low-density lipoprotein receptor–related protein 1 in maintaining the integrity of blood vessels by regulating protease activity, as well as matrix deposition by modulating HtrA1 and connective tissue growth factor. Xiao et al²² demonstrated the potential therapeutic use of the soluble Jagged1 because it not only inhibits proliferation of pulmonary VSMCs but also restores VSMCs phenotype from the dedifferentiated to the differentiated state through interference with Notch-Her2 signaling. Finally, Dalvi et al²³ showed that simultaneous exposure of pulmonary VSMCs to viral proteins and cocaine exacerbates

downregulation of bone morphogenetic protein receptor. Taken together, these studies in the recent issues of *ATVB* have substantially contributed to the progress of our understanding on the important roles of VSMCs in the development of vascular diseases.

Adventitial Tissues and Inflammatory Cells for Vascular Function

Cross talk between cells in the blood vessel wall is crucial for vascular homeostasis and diseases. EC dysfunction leads to the expression of adhesion molecules for inflammatory cells, which promotes the development of vascular diseases. The adhesion and migration of inflammatory cells produce an oxidizing environment, which is critical for the progression of vascular diseases. The accumulating inflammatory cells produce abundant ROS and secrete inflammatory cytokines and growth factors, all of which contribute to EC dysfunction and VSMC proliferation.²⁴ In a recent issue of *ATVB*, Karper et al²⁵ identified a crucial role of toll-like receptor accessory molecule RP105 on circulating inflammatory cells in atherosclerotic plaque formation. In addition, Shiomi et al²⁶ demonstrated that coronary vasospasm could trigger coronary plaque injury and acute ischemic myocardial damage in Watanabe heritable hypercholesterolemic rabbits. Furthermore, Callegari et al²⁷ demonstrated that osteoprotegerin derived from the bone marrow block the progression of vascular calcification.²⁷ About the pathogenesis of aneurysms, Marinkovic et al²⁸ performed an interesting study to limit aneurysm growth by inhibition of inflammation and reducing EC activation with immunosuppressive drug azathioprine. They showed that azathioprine has an anti-inflammatory effect in ECs and inhibits Rac1 and c-Jun-terminal-N-kinase activation, which may explain its protective role in the development of aneurysm formation and severity.²⁸ The N-terminal lectin-like domain of thrombomodulin is known to have an anti-inflammatory function. The recent study by Lin et al²⁹ provided a mechanism showing that recombinant thrombomodulin domain 1 can inhibit inflammation by binding to its carbohydrate ligand Le(y). However, peroxisome proliferator-activated receptor (PPAR)- γ coactivator 1 α is an important mediator of mitochondrial biogenesis and function. Because dysfunctional mitochondria are involved in the pathogenesis of vascular disease, Kröller-Schön et al³⁰ examined the effects of peroxisome PPAR- γ coactivator 1 α deficiency during chronic angiotensin II treatment in vivo. Peroxisome PPAR- γ coactivator 1 α deletion caused vascular dysfunction and inflammation during chronic angiotensin II infusion by increasing mitochondrial ROS production.³⁰ Furthermore, the findings by Doyon et al³¹ provided new insights into the molecular mechanism of the pathological role of angiotensin II and assist in identifying the beneficial effects of I κ B kinase- β inhibition for the treatment of cardiovascular diseases. Finally, perivascular adipose tissue wraps blood vessels and modulates vasoreactivity by secretion of vasoactive molecules. Mammalian target of rapamycin complex 2 has been shown to control inflammation and is expressed in adipose tissue. Bhattacharya et al³² identified mammalian target of rapamycin complex 2 as a critical regulator of perivascular adipose tissue-directed protection of normal vascular tone. They concluded that modulation of

mammalian target of rapamycin complex 2 activity in adipose tissue is a potential therapeutic approach for inflammation-related vascular damage.³²

Summary

In summary, the interactions among ECs, VSMCs, adventitial tissues, and inflammatory cells regulate vascular functions and diseases. As reviewed in this highlights, recent publications in *ATVB* have significantly contributed to our understanding on the roles of each vascular cell component for vascular functions in health and disease. These studies have demonstrated the novel mechanistic roles of secreted factors, as well as intracellular signaling for endothelial homeostasis, VSMC proliferation, and inflammatory cell migration. Indeed, we are getting closer to the understanding of fundamental vascular functions and causative cell signaling pathways, all of which should be useful for the development of novel therapeutic strategies in cardiovascular medicine.

Sources of Funding

This work was supported, in part, by the Grant-in-Aid for Tohoku University Global COE for Conquest of Signal Transduction Diseases with Network Medicine, the Grants-in-Aid for Scientific Research (21790698, 23659408, and 24390193) from the Ministry of Education, Culture, Sports, Science and Technology, Tokyo, Japan, and the Grants-in-Aid for Scientific Research (10102895) from the Ministry of Health, Labor, and Welfare, Tokyo, Japan.

Disclosures

H. Shimokawa is a consultant of Asahi Kasei Pharma. K. Satoh reports no conflicts.

References

1. Walshe TE, dela Paz NG, D'Amore PA. The role of shear-induced transforming growth factor- β signaling in the endothelium. *Arterioscler Thromb Vasc Biol.* 2013;33:2608–2617.
2. Enkhjargal B, Godo S, Sawada A, Suvd N, Saito H, Noda K, Satoh K, Shimokawa H. Endothelial AMP-activated protein kinase regulates blood pressure and coronary flow responses through hyperpolarization mechanism in mice. *Arterioscler Thromb Vasc Biol.* 2014;34:1505–1513.
3. Erbilgin A, Siemers N, Kayne P, Yang WP, Berliner J, Lusis AJ. Gene expression analyses of mouse aortic endothelium in response to atherogenic stimuli. *Arterioscler Thromb Vasc Biol.* 2013;33:2509–2517.
4. Choi S, Kim JA, Na HY, Cho SE, Park S, Jung SC, Suh SH. Globotriaosylceramide induces lysosomal degradation of endothelial KCa3.1 in fabry disease. *Arterioscler Thromb Vasc Biol.* 2014;34:81–89.
5. Satoh K. Globotriaosylceramide induces endothelial dysfunction in fabry disease. *Arterioscler Thromb Vasc Biol.* 2014;34:2–4.
6. Wood KC, Cortese-Krott MM, Kovacic JC, Noguchi A, Liu VB, Wang X, Raghavachari N, Boehm M, Kato GJ, Kelm M, Gladwin MT. Circulating blood endothelial nitric oxide synthase contributes to the regulation of systemic blood pressure and nitrite homeostasis. *Arterioscler Thromb Vasc Biol.* 2013;33:1861–1871.
7. Ciccarelli M, Sorriento D, Franco A, Fusco A, Del Giudice C, Annunziata R, Cipolletta E, Monti MG, Dorn GW II, Trimarco B, Iaccarino G. Endothelial G protein-coupled receptor kinase 2 regulates vascular homeostasis through the control of free radical oxygen species. *Arterioscler Thromb Vasc Biol.* 2013;33:2415–2424.
8. Zippel N, Malik RA, Frömel T, Popp R, Bess E, Strilic B, Wettschreck N, Fleming I, Fisslthaler B. Transforming growth factor- β -activated kinase 1 regulates angiogenesis via AMP-activated protein kinase- α 1 and redox balance in endothelial cells. *Arterioscler Thromb Vasc Biol.* 2013;33:2792–2799.
9. Iso T, Maeda K, Hanaoka H, et al. Capillary endothelial fatty acid binding proteins 4 and 5 play a critical role in fatty acid uptake in heart and skeletal muscle. *Arterioscler Thromb Vasc Biol.* 2013;33:2549–2557.

10. Wagner NM, Bierhansl L, Nöldge-Schomburg G, Vollmar B, Roesner JP. Toll-like receptor 2-blocking antibodies promote angiogenesis and induce ERK1/2 and AKT signaling via CXCR4 in endothelial cells. *Arterioscler Thromb Vasc Biol*. 2013;33:1943–1951.
11. Tatematsu S, Francis SA, Natarajan P, Rader DJ, Saghatelian A, Brown JD, Michel T, Plutzky J. Endothelial lipase is a critical determinant of high-density lipoprotein-stimulated sphingosine 1-phosphate-dependent signaling in vascular endothelium. *Arterioscler Thromb Vasc Biol*. 2013;33:1788–1794.
12. Parastatidis I, Weiss D, Joseph G, Taylor WR. Overexpression of catalase in vascular smooth muscle cells prevents the formation of abdominal aortic aneurysms. *Arterioscler Thromb Vasc Biol*. 2013;33:2389–2396.
13. Soe NN, Sowden M, Baskaran P, Smolock EM, Kim Y, Nigro P, Berk BC. Cyclophilin A is required for angiotensin II-induced p47phox translocation to caveolae in vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol*. 2013;33:2147–2153.
14. Shimizu T, Fukumoto Y, Tanaka S, Satoh K, Ikeda S, Shimokawa H. Crucial role of ROCK2 in vascular smooth muscle cells for hypoxia-induced pulmonary hypertension in mice. *Arterioscler Thromb Vasc Biol*. 2013;33:2780–2791.
15. Ikeda S, Satoh K, Kikuchi N, Miyata S, Suzuki K, Omura J, Shimizu T, Kobayashi K, Kobayashi K, Fukumoto Y, Sakata Y, Shimokawa H. Crucial role of Rho-kinase in pressure overload-induced right ventricular hypertrophy and dysfunction in mice. *Arterioscler Thromb Vasc Biol*. 2014;34:1260–1271.
16. Xiao Q, Zhang F, Grassia G, Hu Y, Zhang Z, Xing Q, Yin X, Maddaluno M, Drung B, Schmidt B, Maffia P, Ialenti A, Mayr M, Xu Q, Ye S. Matrix metalloproteinase-8 promotes vascular smooth muscle cell proliferation and neointima formation. *Arterioscler Thromb Vasc Biol*. 2014;34:90–98.
17. Turlo KA, Scapa J, Bagher P, Jones AW, Feil R, Korhuis RJ, Segal SS, Iruela-Arispe ML. β 1-integrin is essential for vasoregulation and smooth muscle survival in vivo. *Arterioscler Thromb Vasc Biol*. 2013;33:2325–2335.
18. Sutliff RL, Hilenski LL, Amanso AM, Parastatidis I, Dikalova AE, Hansen L, Datla SR, Long JS, El-Ali AM, Joseph G, Gleason RL, Jr, Taylor WR, Hart CM, Griendling KK, Lassègue B. Polymerase delta interacting protein 2 sustains vascular structure and function. *Arterioscler Thromb Vasc Biol*. 2013;33:2154–2161.
19. Song P, Zhou Y, Coughlan KA, Dai X, Xu H, Viollet B, Zou MH. Adenosine monophosphate-activated protein kinase- α 2 deficiency promotes vascular smooth muscle cell migration via S-phase kinase-associated protein 2 upregulation and E-cadherin downregulation. *Arterioscler Thromb Vasc Biol*. 2013;33:2800–2809.
20. Revuelta-López E, Castellano J, Roura S, Gálvez-Montón C, Nasarre L, Benitez S, Bayes-Genis A, Badimon L, Llorente-Cortés V. Hypoxia induces metalloproteinase-9 activation and human vascular smooth muscle cell migration through low-density lipoprotein receptor-related protein 1-mediated Pyk2 phosphorylation. *Arterioscler Thromb Vasc Biol*. 2013;33:2877–2887.
21. Muratoglu SC, Belgrave S, Hampton B, Migliorini M, Coksaygan T, Chen L, Mikhailenko I, Strickland DK. LRP1 protects the vasculature by regulating levels of connective tissue growth factor and HtrA1. *Arterioscler Thromb Vasc Biol*. 2013;33:2137–2146.
22. Xiao Y, Gong D, Wang W. Soluble JAGGED1 inhibits pulmonary hypertension by attenuating notch signaling. *Arterioscler Thromb Vasc Biol*. 2013;33:2733–2739.
23. Dalvi P, O'Brien-Ladner A, Dhillon NK. Downregulation of bone morphogenetic protein receptor axis during HIV-1 and cocaine-mediated pulmonary smooth muscle hyperplasia: implications for HIV-related pulmonary arterial hypertension. *Arterioscler Thromb Vasc Biol*. 2013;33:2585–2595.
24. Tedgui A, Ait-Oufella H. Innate lymphoid cells: new players in atherosclerosis? *Arterioscler Thromb Vasc Biol*. 2013;33:2697–2698.
25. Karper JC, de Jager SC, Ewing MM, de Vries MR, Bot I, van Santbrink PJ, Redeker A, Mallat Z, Binder CJ, Arens R, Jukema JW, Kuiper J, Quax PH. An unexpected intriguing effect of Toll-like receptor regulator RP105 (CD180) on atherosclerosis formation with alterations on B-cell activation. *Arterioscler Thromb Vasc Biol*. 2013;33:2810–2817.
26. Shiomi M, Ishida T, Kobayashi T, Nitta N, Sonoda A, Yamada S, Koike T, Kuniyoshi N, Murata K, Hirata K, Ito T, Libby P. Vasospasm of atherosclerotic coronary arteries precipitates acute ischemic myocardial damage in myocardial infarction-prone strain of the Watanabe heritable hyperlipidemic rabbits. *Arterioscler Thromb Vasc Biol*. 2013;33:2518–2523.
27. Callegari A, Coons ML, Ricks JL, Yang HL, Gross TS, Huber P, Rosenfeld ME, Scatena M. Bone marrow- or vessel wall-derived osteoprotegerin is sufficient to reduce atherosclerotic lesion size and vascular calcification. *Arterioscler Thromb Vasc Biol*. 2013;33:2491–2500.
28. Marinkovic G, Hibender S, Hoogenboezem M, van Broekhoven A, Girigorie AF, Bleeker N, Hamers AA, Stap J, van Buul JD, de Vries CJ, de Waard V. Immunosuppressive drug azathioprine reduces aneurysm progression through inhibition of Rac1 and c-Jun-terminal-N-kinase in endothelial cells. *Arterioscler Thromb Vasc Biol*. 2013;33:2380–2388.
29. Lin WL, Chang CF, Shi CS, Shi GY, Wu HL. Recombinant lectin-like domain of thrombomodulin suppresses vascular inflammation by reducing leukocyte recruitment via interacting with Lewis Y on endothelial cells. *Arterioscler Thromb Vasc Biol*. 2013;33:2366–2373.
30. Kröll-Schön S, Jansen T, Schüler A, Oelze M, Wenzel P, Hausding M, Kerahrodi JG, Beisele M, Lackner KJ, Daiber A, Münzel T, Schulz E. Peroxisome proliferator-activated receptor γ , coactivator 1 α deletion induces angiotensin II-associated vascular dysfunction by increasing mitochondrial oxidative stress and vascular inflammation. *Arterioscler Thromb Vasc Biol*. 2013;33:1928–1935.
31. Doyon P, van Zuylen WJ, Servant MJ. Role of I κ B kinase- β in the growth-promoting effects of angiotensin II *in vitro* and *in vivo*. *Arterioscler Thromb Vasc Biol*. 2013;33:2850–2857.
32. Bhattacharya I, Dräger K, Albert V, Contassot E, Damjanovic M, Hagiwara A, Zimmerli L, Humar R, Hall MN, Battagay EJ, Haas E. Rictor in perivascular adipose tissue controls vascular function by regulating inflammatory molecule expression. *Arterioscler Thromb Vasc Biol*. 2013;33:2105–2111.

KEY WORDS: endothelial cells ■ muscle, smooth, vascular